

# Precipitation variability over the East African region and its relation with circulation patterns

N. Souverijns<sup>1</sup>, W. Thiery<sup>1</sup>, M. Demuzere<sup>1</sup>, N.P.M. van Lipzig<sup>1</sup>

<sup>1</sup> EES, University of Leuven, Belgium, [niels.souverijns@student.kuleuven.be](mailto:niels.souverijns@student.kuleuven.be)

## 1. Motivation




Figure 1. Study area weighed normalised PDSI index (data: KNMI)

- High interannual precipitation variability poses severe risks for population and food security



Figure 2. Live Aid Concert 1985

## 2. Setup




Figure 3. Study area

- 1981-2010
- ERA Interim data
- Relate Mean Sea Level Pressure to:
  - Total Precipitation
  - Large-Scale Precipitation
  - Convective Precipitation

## 3. COST733Class

- Classify Mean Sea Level Pressure in different patterns
- Evaluate different classification algorithms with different separability measures
- Separating pressure patterns is considered most important
- Neural-network classifications perform best in separating pressure patterns




Figure 4. Separability analysis regarding Mean Sea Level Pressure




Figure 5. Separability analysis regarding precipitation

## 4. Weather Atlas




Figure 6. Results of the best classification algorithm

Month	C1	C2	C3	C4	C5	C6	C7	C8	C9
January	0	45.7	0	4.6	0	8.1	0	41.6	0
February	0	42.7	0.1	9.2	0	24.3	0	23.7	0
March	0	20.6	5.4	18.2	0.1	45.5	0	10.2	0
April	0	2.6	52.5	11.2	2.6	28.8	0	2.3	0
May	5.4	0	52.4	1.0	14.4	2.8	3.0	0	21.0
June	7.0	0	2.2	0	0.7	0	33.4	0	56.7
July	2.5	0	0	0	0.1	0	56.8	0	40.6
August	14.7	0	0	0	4.3	0	47.2	0	33.8
September	48.6	0	0.4	1.4	32.8	0.1	7.8	0	8.9
October	9.1	0	1.7	49	35.3	3.3	0	1.6	0
November	0	9.4	0.6	51.4	3.2	10.3	0	25.1	0
December	0	36.4	0	13.3	0	6.3	0	44.0	0

Figure 7. Occurrence of different circulation types per month in the period 1981-2010

- Distinct seasonal patterns are visible
- Precipitation patterns are clearly related to circulation
- Droughts / wet spells are mostly related to differences in interannual frequency of circulation patterns

## 5. (Near-) future results

- Classify pressure patterns of COSMO-CLM present (1989-2010) and future (2081-2100) simulations
- RCP8.5 scenario
- Relate changes in precipitation and PDSI (variability) to changes in circulation patterns
- Changes in circulation pattern frequency and the patterns itself
- Relate circulation pattern occurrence to El Niño / La Niña events
- Possible relation with dry / wet periods
- Clear correlation between Nino3.4 index and precipitation observations in the long rain season (October, November, December).
- Possible lagged relation in the short rain season (March, April, May)




Figure 8. Nino 3.4 index (data: KNMI)




Figure 9. Correlation between Nino 3.4 and precipitation (CRU) for Oct, Nov and Dec (data: KNMI)

